


Recombinant Human CD26 (N-Fc)

Catalog#:P01824 Derived from Human Cells

DESCRIPTION	<p>Recombinant Human Dipeptidyl Peptidase 4 is produced by our Mammalian expression system and the target gene encoding Asn29-Pro766 is expressed with a Fc tag at the N-terminus.</p> <p>Accession#: P27487</p> <p>Known as: Dipeptidyl peptidase 4; ADABP; Adenosine deaminase complexing protein 2; ADCP-2; Dipeptidyl peptidase IV; DPP IV; T-cell activation antigen CD26</p>
FORMULATION	Supplied as a 0.2 µm filtered solution of PBS, pH 7.4.
SHIPPING	<p>The product is shipped on dry ice/polar packs.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Store at ≤-70°C, stable for 6 months after receipt.</p> <p>Store at ≤-70°C, stable for 3 months under sterile conditions after opening.</p> <p>Please minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass: 111.7kDa AP Mol Mass: 105-130kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test.</p>
BACKGROUND	<p>CD26 is a signal-anchor for type II membrane protein that belongs to the peptidase S9B family. CD26 is expressed specifically in lymphatic vessels but not in blood vessels in the skin, small intestine, esophagus, ovary, breast and prostate glands. It acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. It's binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappaB activation in a T-cell receptor/CD3-dependent manner. Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. It may be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, it enhanced cell proliferation, a process inhibited by GPC3. It acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones.</p>
<p style="text-align: center;">SDS-PAGE</p>  <p style="text-align: center;">kDa MK R</p> <p style="text-align: center;">170 130 95 72 55 43 34</p>	