



5×蛋白上样缓冲液（含巯基还原剂）

货号：P1041

规格：10 mL

保存：-20°C保存，建议分装冻存，避免反复冻融。有效期 12 个月。

产品简介：

本产品适用于 SDS-PAGE(SDS 变性聚丙烯酰胺凝胶电泳)时作蛋白质上样用。其主要成份为 SDS，巯基还原剂，溴酚蓝，缓冲盐溶液等。

SDS 可与蛋白质结合使蛋白质—SDS 复合物上带有大量的负电荷，这时蛋白质本身的电荷完全被 SDS 掩盖，消除了各种蛋白质本身电荷的差异，SDS 还可以断开分子内和分子间的氢键，破坏蛋白质分子的二级和三级结构， β -巯基乙醇可以断开半胱氨酸残基之间的二硫键，破坏蛋白质的四级结构，消除了蛋白结构之间的差异。最终无电荷及结构上差异的蛋白（亚单位），电泳速度只是与其分子量大小有关。溴酚蓝用作电泳时的指示剂，可大概指示电泳结束的时间。

使用说明（仅供参考）：

1. 请按每 40 μ L 蛋白样品加入 10 μ L 上样缓冲液的比例（5 倍稀释）来使用。如果蛋白浓度过高，可用双蒸水稀释。
2. 混匀后，100°C水浴加热 3-5 分钟，使蛋白变性。
3. 冷却到室温后，10000-14000 rpm 离心 2-5 分钟，取上清直接上样电泳即可。

注意事项：

1. 聚丙烯酰胺凝胶浓度为 8%时溴酚蓝指示条带的位置大概在 30kd 左右，胶浓度 12%时，约在 20kd 左右，胶浓度 15%时，大概在 10kd。请根据自己目标条带来判断结果电泳时间。
2. 本试剂因含巯基还原剂，有一定的毒性，为了您的安全和健康，请穿实验服并戴一次性手套操作。

相关产品：

P1015	4×蛋白上样缓冲液（含 DTT）
P1017	4×非变性蛋白上样缓冲液
P0012	10×丽春红染色液
P1300	考马斯亮蓝快速染色液
P1200	SDS-PAGE 凝胶制备试剂盒
PR1700	预染次高分子量蛋白 Marker

相关文献：

- [1] Zhihui Wang,Zhiquan Zhu,Zhong Lin,et al. miR-429 suppresses cell proliferation, migration and invasion in nasop -haryngeal carcinoma by downregulation of TLN1. Cancer Cell International. April 2019. (IF 3.439)
- [2] Youbing Tu,Yubing Liang,Yong Xiao,et al. Dexmedetomidine attenuates the neurotoxicity of propofol toward primary hippocampal neurons in vitro via Erk1/2/CREB/BDNF signaling pathways. Drug Design, Development and Therapy. February 2019. (IF 3.486)

注：更多使用本产品的文献请参考索莱宝官网。