


Recombinant Mouse IL-4RA

Catalog#:P00919 Derived from Human Cells

DESCRIPTION	<p>Recombinant Mouse Interleukin-4 Receptor Subunit Alpha is produced by our Mammalian expression system and the target gene encoding Ile26-Arg233 is expressed with a human IgG1 Fc tag at the C-terminus.</p> <p>Accession#:P16382</p> <p>Known as: Interleukin-4 receptor subunit alpha; IL-4R-alpha; CD124; IL4-BP; Soluble IL-4R-alpha.</p>
FORMULATION	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt.</p> <p>Reconstituted protein solution can be stored at 2-8$^{\circ}\text{C}$ for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100$\mu\text{g}/\text{ml}$.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass:51.5kDa AP Mol Mass:65-80kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1ng/μg (1 EU/μg) as determined by LAL test.</p>
BACKGROUND	<p>Interleukin-4 receptor subunit alpha(IL-4RA), also known as Soluble IL-4 receptor subunit alpha, belongs to the type I cytokine receptor family and type 4 subfamily. It expressed in both Th1 and Th2 cells. It functions as receptor for both interleukin 4 and interleukin 13 and couples to the JAK1/2/3-STAT6 pathway. The IL4 response is involved in promoting Th2 differentiation. The IL4/IL13 responses are involved in regulating IgE production and chemokine and mucus production at sites of allergic inflammation. In certain cell types, IL-4RA can signal through activation of insulin receptor substrates, IRS1/IRS2. The functional IL4 receptor is formed by initial binding of IL4 to IL4R. Subsequently it recruits to the complex of the common gamma chain. In immune cells, IL-4RA creates a type I receptor. In non-immune cells, it forms a type II receptor with of IL13RA1. IL4R can also interact with the IL13/IL13RA1 complex to form a similar type II receptor and interacts with the SH2-containing phosphatases, PTPN6/SHIP1, PTPN11/SHIP2 and INPP5D/SHIP.</p>
SDS-PAGE	 <p>kDa MK R</p> <p>120</p> <p>90</p> <p>60</p> <p>40</p> <p>30</p> <p>20</p>