

## Avidin

**Cat:** A8280

**Specification:** 1mg

**Storage:** Store at -20°C, and it is valid for 6 years.

### Product Information

**CAS:** 1405-69-2

**English name:** Avidin

**Appearance (Character):** White powder

**Grade:** Reagent Grade

**Purity:** ≥95.0%

**Source:** Chicken egg white\_

**Solubility:** 10mg/ml in Water; Soluble in 0.1M phosphate buffer pH6.8, reference concentration 1mg/ml.

### Introduction

Avidin is a tetrameric glycoprotein with an approximate molecular mass of 66-67 kDa. It is composed of four subunits with each subunit containing 128 identical amino acid residues and a variable carbohydrate moiety. The subunits may vary slightly in molecular mass due to the carbohydrate composition. Only Asp is glycosylated. The carbohydrate moiety can have at least three different carbohydrate structural types. Avidin can be dissociated into subunits under strongly denaturing conditions. Each subunit is separately capable of binding biotin with a dissociation constant ( $K_d$ ) =  $10^{-7}M$ .

The avidin-biotin association constant ( $K_a = 10^{15} M^{-1}$ ) is one of the strongest affinities known. The complex is stable to 100°C, and significantly stable to detergents and denaturants. Avidin is stable to 85°C without biotin. Biophysical studies of the avidin-biotin complex have implicated particular tryptophan and lysine residues in the biotin-binding site of avidin. The crystal structure of avidin has been published. The crystal structure of a deglycosylated form of avidin complexed with biotin has been reported. Because of the stability of both avidin and biotin, each of these molecules has been used as "labels" for antibodies, fluorescent dyes, proteins, and other molecules of interest to biochemists. Avidin and biotin have each been incorporated into immobilized matrices. (The only way that monomeric avidin can exist is through its attachment to an agarose support.)

This product is purified using affinity chromatography. It is dialyzed extensively against deionized water before being lyophilized. The basic procedures are modified from literature preparations. This product is sold by protein content (determined by E1%280).

**Unit Definition:** One unit will bind 1.0 µg of d-biotin.

**Unit activity:** ≥10 units per mg protein

**Isoelectric point (pI):** 10

**Kd for the avidin-biotin complex:** 10<sup>-15</sup> M (neutral pH)

**Binding capacity:**

For tetrameric avidin, the theoretical maximum is 4 moles biotin:1 mole avidin or 15 µg biotin/mg protein.

**Reported fluorescent wavelength:**

338 nm (avidin)

328 nm (avidin-biotin complex)

**Preparation Instructions**

Avidin is very soluble both in water, up to 20 mg/mL, and in salt solutions. Avidin solutions are stable over a wide range of pH and temperatures, particularly when combined with biotin. Avidin can be crystallized from ammonium sulfate at  $> 2.5$  M at pH 5. Since one tryptophan residue per subunit is involved in the binding site, avidin can be inactivated by oxidizing agents such as ozone, peroxide, or strong light. Solutions should be stored at  $-20^{\circ}\text{C}$ .

**Storage/Stability**

The avidin-biotin complex is even more heat stable than avidin alone. It is only 10% dissociated after 15 minutes at  $100^{\circ}\text{C}$ . It is not completely dissociated after 60 minutes at  $100^{\circ}\text{C}$ . The complex can be quantitatively dissociated only under autoclaving conditions, e.g.  $120^{\circ}\text{C}$ , 15 minutes. When avidin was reduced in the presence of 9 M urea, its biotin-binding activity was unchanged. The protein was denatured and lost biotin-binding activity as the pH was gradually lowered to pH 1. However, when the pH was raised to pH 3, avidin regained native configuration and binding activity. The complex is also extremely stable at high pH, being only 20% ionized even at pH 13.