

Creatinine Content Assay Kit (Creatine Oxidase Method)

Note: It is necessary to predict 2-3 large difference samples before the formal determination.

Operation Equipment: Spectrophotometer/Microplate reader

Cat No: BC4915

Size: 100T/96S

Components:

Extracting solution I: Liquid 110 mL×1. Storage at 4°C.

Extracting solution II: Liquid 20 mL×1. Storage at 4°C.

Reagent I: Powder ×2. Storage at -20°C. Before use, add 1.1 mL of distilled water to each tube and dissolve it completely. The unused reagents is divided and then stored at -20°C in two weeks.

Reagent II: Powder ×2. Storage at -20°C. Before use, add 1.05 mL of distilled water to each tube and dissolve it completely. The unused reagents is divided and then stored at -20°C in two weeks.

Reagent III: Powder ×2. Storage at -20°C. Before use, add 1 mL of distilled water to each tube (100T/96S) and dissolve it completely. For the convenience of storage, give one more. The unused reagents is divided and then stored at -20°C in two weeks.

Reagent III working solution: according to the required amount of the test, according to the ratio of Reagent III: distilled water = 1:9, fully mix and use now.

Reagent IV: Powder ×1. Storage at 4°C. Before use, add 1 mL of distilled water and dissolve it completely. The unused reagents is divided and then stored at -20°C.

Reagent V: Powder ×2. Storage at -20°C. Before use, add 2.2 mL of distilled water to each tube and dissolve it completely. The unused reagents is divided and then stored at -20°C.

Reagent VI A: Liquid 5 mL ×2. Storage at 4°C.

Reagent VI B: Liquid 5 mL ×2. Storage at 4°C. Before use, according to the amount required by the experiment, the mixture shall be fully mixed according to the ratio of Reagent VIA: Reagent VIB = 1:1, and prepare when the solution will be used.

Standard solution: Powder ×1, 1 mg of creatinine. Before use, add 1 mL of distilled water to fully dissolve, i.e. 1 mg/mL standard solution. Before use, dilute it to 200 µg/mL as the standard solution.

Product Description:

Creatinine (Cre), the chemical formula is $C_4H_7N_3O$, is a product of muscle metabolism in the human body, which is mainly excreted by glomerular filtration. Serum creatinine comes from both exogenous and endogenous sources. Exogenous creatinine is the product of meat metabolism in the body; endogenous creatinine is the product of muscle metabolism in the body.

The creatinine in the sample is hydrolyzed to creatine under the catalysis of creatinine enzyme. Under the catalysis of creatinase, creatine hydrolyzes to creatine and urea. Sarcosine is oxidized to hydrogen peroxide under the catalysis of sarcosine oxidase. Peroxidase catalyzed the oxidation of 4-aminoantipyrine coupling phenol by hydrogen peroxide to form colored compounds with

characteristic absorption peak at 505nm.

Reagents and Equipment Required but Not Provided:

Spectrophotometer/microplate reader, low temperature centrifuge, transferpettor, micro glass cuvette/96 well flat-bottom plate, mortar/homogenizer, ice and distilled water, ultrasonic crusher.

Procedure:

I. Sample preparation (the sample size can be adjusted appropriately, and the specific proportion can be referred to the literature):

1. Preparation of bacteria and cell samples: according to the cell number (10^4): the volume of Extracting solution I (mL) is 500 ~ 1000:1 (it is recommended to add 1 mL Extracting solution I to 5 million cells), ice bath ultrasonic wave is used to crush cells (power 300W, ultrasonic 3 seconds, interval 9 seconds, total time 5 min); centrifugation at 4°C, 12000 g for 10 min, take 0.8 mL of supernatant, and then add 0.15 mL of Extracting solution II, After centrifugation at 4°C and 12000 g for 10 min, the supernatant is taken for determination.

2. Preparation of tissue samples: according to the ratio of mass (g): the volume of Extracting solution I (mL) of 1:5~10 (it is recommended to weigh about 0.1 g of tissue and add 1 mL of Extracting solution I), add Extracting solution I, homogenize in ice bath, centrifuge at 4°C, 12000g for 10 min, take 0.8 mL of supernatant, and then add 0.15 mL of Extracting solution II, centrifuge at 4°C, 12000g for 10 min, take supernatant for testing.

3. Serum (plasma): take 100 μ L of serum(plasma) and add 1 mL of Extracting solution I, centrifuge at 4°C, 12000g for 10 min, take 0.8 mL of supernatant, then add 0.15 mL of Extracting solution II, centrifuge at 4°C, 12000g for 10 min, and then take the supernatant for testing.

II. Determination procedure:

1. Preheat the Spectrophotometer/microplate reader for 30 minutes, adjust the wavelength to 505 nm, set zero with distilled water.

2. Add reagents with the following list:

Reagent (μ L)	Test tube (T)	Blank tube (B)	Standard (S)
Sample	20	-	-
Distilled water	-	20	-
Standard solution	-	-	20
Reagent I	20	20	20
Reagent II	20	20	20
Reagent III working solution	5	5	5
Reagent IV	5	5	5
Mix well and react for 10 min at 37°C (mammalian) or 25°C (other species).			
Reagent IV	40	40	40
Reagent V	90	90	90

Mix well, color for 30 min at 37°C (mammalian) or 25°C (other species). The absorbance at 505 nm is determined. They are respectively recorded as A_T , A_B and A_S . $\Delta A_T = A_T - A_B$, $\Delta A_S = A_S - A_B$.

Note: the Blank tube only needs 1-2 times.

III. Calculation:

1. Calculation formula

(1) Calculated according to protein concentration

$$\text{Creatinine content } (\mu\text{g}/\text{mg prot}) = C_S \times \Delta A_T \div \Delta A_S \times V_S \div (V_S \times \text{Cpr}) = 200 \times \Delta A_T \div \Delta A_S \div \text{Cpr}$$

(2) Calculated by sample quality

$$\text{Creatinine content } (\mu\text{g}/\text{g mass}) = C_S \times \Delta A_T \div \Delta A_S \times (V_U + V_{E2}) \div (W \times V_U \div V_{E1}) = 237.5 \times \Delta A_T \div \Delta A_S \div W$$

(3) Calculated by the number of bacteria or cells

$$\text{Creatinine content } (\mu\text{g}/10^4 \text{ cells}) = C_S \times \Delta A_T \div \Delta A_S \times (V_U + V_{E2}) \div (\text{cell number} \times V_S \div V_{E1}) = 237.5 \times \Delta A_T \div \Delta A_S \div \text{cell number}$$

(4) Calculated according to the volume of serum

$$\text{Creatinine content } (\mu\text{g}/\text{mL}) = C_S \times \Delta A_T \div \Delta A_S \times (V_U + V_{E2}) \div (V_{E1} + V_L) = 2612.5 \times \Delta A_T \div \Delta A_S$$

C_S : standard tube concentration, 200 $\mu\text{g}/\text{mL}$; V_S : add sample volume, 20 $\mu\text{L} = 0.02 \text{ mL}$; V_U : volume of supernatant during extraction, 0.8 mL; V_{E1} : add the volume of Extracting solution I, 1 mL; V_{E2} : add the volume of Extracting solution II, 0.15 mL; W : sample mass, g; Cpr : sample protein concentration, mg/mL ; cell number: 10^4 ; V_L : the volume of liquid sample, 0.1 mL.

Note:

1. The supernatant can not be used for the determination of protein concentration. If you want to calculate creatinine content with protein concentration, you need to take another tissue or serum (plasma), that is, take the same mass (volume) of tissue (serum (plasma)) with 1.1875 mL PBS (normal saline) homogenate (equivalent to the final sample supernatant of the extraction step), and use BCA method to determine protein concentration.

2. If the test tube absorbance value exceeds that of the standard tube, it is recommended to dilute the sample before measuring. If the absorbance value is too small, it is recommended to increase the sample before determination.

Experimental examples:

1. Take 0.1 g of mouse muscle and add 1 mL of Extracting solution I for homogenization and centrifugation. Take 0.8 mL of supernatant and add 0.15 mL Extracting solution II. After centrifugation, operate according to the determination steps. After determination with 96 well plate, calculate $\Delta A_T = A_T - A_B = 0.199 - 0.054 = 0.145$, $\Delta A_S = A_S - A_B = 0.483 - 0.054 = 0.429$. The content is calculated according to the sample mass.

The content of creatinine ($\mu\text{g}/\text{g mass}$) = $237.5 \times \Delta A_T \div \Delta A_S \div W = 802.7 \mu\text{g}/\text{g mass}$.

2. Take 100 μL of bovine serum, add 1 mL of Extracting solution I, take 0.8 mL of supernatant and add 0.15 mL of Extracting solution II, the centrifugal supernatant, and then operate according to the determination steps. After determination with 96 well plate, calculate $\Delta A_T = A_T - A_B = 0.138 - 0.054 = 0.084$, $\Delta A_S = A_S - A_B = 0.483 - 0.054 = 0.429$. The content is calculated according to the volume of

liquid.

The content of creatinine ($\mu\text{g/mL}$) = $2612.5 \times \Delta A_T \div \Delta A_S = 511.5 \mu\text{g/mL}$ serum.