

Arginase Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Catalog Number: BC5550

Size: 50T/24S

Components:

Extract solution I: Liquid 30ml×1, store at 2-8°C.

Extract solution II: Liquid 0.3ml×1, store at -20°C.

Extract solution: According to the number of samples, prepare in the ratio of Extract I: Extract II = 990μL: 10μL (1T). It is forbidden to add all of Extract II to Extract I at once.

Reagent I: Powder×1, store at 2-8°C. Dissolve well with 9.6mL of reagent II before use. Unused reagents can be stored at 2-8°C for up to 4 weeks. Do not store at -20°C.

Reagent II: Liquid 10ml×1, store at 2-8°C.

Reagent III: Liquid 25ml×1, store at 2-8°C.

Reagent IV: Liquid 36ml×1, store at 2-8°C.

Reagent V: Liquid 15ml×1, store at 2-8°C.

Standard: Liquid 1ml×1, store at 2-8°C. 1 mol/L (1000 μmol/mL) urea standard solution.

Product Description:

Arginase, also known as L-arginine urea hydrolase or L-arginine amidinyl hydrolase, is a manganese metalloenzyme. Arginase is found in bacteria, yeast, plants, invertebrates and vertebrates, and is thought to have first appeared in bacteria. The main function of arginase in microorganisms is probably to participate in maintaining the dynamic balance of L-arginine and in regulating a variety of important metabolic processes.

Arginase catabolic L-arginine into L-ornithine and urea, urea reacts with α-isonitrosopropiophenone to produce a derivative with an absorption peak at 560 nm. By measuring the production of urea, the size of the arginine activity can be calculated.

Reagents and Equipment Required but Not Provided:

Spectrophotometer, low temperature centrifuge, adjustable transferpettor, water bath/ constant temperature incubator, 1ml glass cuvette, mortar/homogenizer/cell ultrasonicator, ice and distilled water.

I. Sample extraction:

1. Tissue:

Accordance the ratio of tissue(g) : extract solution volume (mL)=1: 5~10 (add 1 mL of extract solution to 0.1 g of tissue), homogenate on ice. Centrifuge at 12000g for 10 minutes at 4°C, take the supernatant and place it on ice for testing.

2. Bacteria or cells:

Accordance the ratio of cells amount(10^6) : extract solution volume (mL)=5~10: 1 (add 1 mL of extract solution to 5 million cells). Ultrasonic on ice bath to smash cells, (powder 200w, ultrasonic 3s, interval 10s

for 5 minutes). Centrifuge at 12000g for 10 minutes at 4°C, take the supernatant and place it on ice for testing.

II. Determination procedure:

1. Preheat spectrophotometer for 30min, adjust the wavelength to 560 nm, set the counter to zero with distilled water.
2. Preparation of standard solutions: The standards were diluted with distilled water to 50, 25, 12.5, 6.25 and 3.125 $\mu\text{mol/mL}$ respectively.
3. Add the following reagents: (In 1.5m LEP tubes follow the steps in the table below)

Reagent name (μL)	Test tube (T)	Contrast tube (C)	Standard tube (S)	Blank tube (B)
Sample	240	240	-	-
Standard	-	-	240	-
Distilled water	-	-	-	240
Reagent I	120	120	120	120
Reagent III	360	360	360	360
Reagent IV	-	480	480	480
30min at 37°C protected from light				
Reagent IV	480	-	-	-
Centrifuge at 8000g for 5min at room temperature, take another 1.5mLEP and aspirate 1000 μL of supernatant into an EP tube				
Supernatant	1000	1000	1000	1000
Reagent V	200	200	200	200
The reaction was boiled in boiling water and protected from light for 40min. The supernatant was centrifuged at 8000g for 5min at room temperature and the absorbance at 560nm was measured in a cuvette and recorded as A_t , A_c , A_s and A_b . $\Delta A_t = A_t - A_c$, $\Delta A_s = A_s - A_b$. (Standard and blank tubes should only be done 1-2 times, one control tube per assay tube is required.)				

III. Calculation of Arginase activity:

1 Make standard curve:

A standard curve was established based on the concentration of the standard tube (x , $\mu\text{mol/mL}$) and the absorbance ΔA_s (y , ΔA_s). Based on the standard curve, the ΔA_t was substituted into the equation to obtain x ($\mu\text{mol/mL}$).

2 Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalytic production of 1 μmol urea per min every. mg tissue protein

$$\text{Arginase activity (U/mg prot)} = x \times V_s \div (C_{pr} \times V_s) \div T \times F = x \div 30 \div C_{pr} \times F$$

3 Sample weight:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalytic production of 1 μmol urea per min every gram tissue weight.

$$\text{Arginase activity (U/g weight)} = x \times V_s \div (W \div V_e \times V_s) \div T \times F = x \div 30 \div W \times F$$

4 Cells or bacteria:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalytic production of 1 μmol urea per min every 10^6 cells.

$$\text{Arginase activity (U/10}^6 \text{ cell)} = x \times V_s \div (N \div V_e \times V_s) \div T \times F = x \div 30 \div N \times F$$

V_s : Volume of sample added to the reaction system, 0.24mL; V_e : Volume of extraction solution added, 1mL; T: Reaction time, 30min; Cpr: Protein concentration, mg/mL; W: Sample quality, g; N: Number of cells or bacteria, In units of 10^6 ; F: Sample dilution.

Note:

1. If the measured absorbance value is greater than 1.5 or ΔA_t is greater than 1, the sample can be diluted or the first step of the 37°C reaction time can be shortened; if the measured absorbance value or ΔA_t is too small, the sample size can be increased or the first step of the 37°C reaction time can be extended. Simultaneously modify the formulae for the final calculation..

Experimental Examples:

1. Weigh 0.1198g of mouse kidney tissue, add the extract for ice bath homogenization, operate according to the assay procedure, measure with a 1mL glass cuvette to calculate $\Delta A_t = A_t - A_c = 0.740 - 0.033 = 0.707$, bring into the standard curve $y = 0.0225x - 0.0295$ ($R^2 = 0.9966$), $x = 32.733 \mu\text{mol/mL}$, brought into the equation to calculate

$$\text{Arginase activity (U/g mass)} = x \div 30 \div W \times F = 9.108 \text{U/g mass}$$

2. Weigh 0.0989 g of carrot, add the extract to the ice bath and homogenize, follow the assay procedure and measure with a 1 mL glass cuvette to calculate $\Delta A_t = A_t - A_c = 0.109 - 0.013 = 0.096$, bring into the standard curve $y = 0.0225x - 0.0295$ ($R^2 = 0.9966$), $x = 5.578 \mu\text{mol/mL}$, and Brought into the equation to calculate:

$$\text{Arginase activity (U/g mass)} = x \div 30 \div W \times F = 1.880 \text{U/g mass}$$

References:

[1] Chen H, Mccaig B C, Melotto M, et al. Regulation of Plant Arginase by Wounding, Jasmonate, and the Phytotoxin Coronatine[J]. Journal of Biological Chemistry, 2004, 279(44):45998-46007.

[2] Ishii N, Ikenaga H, Carmines P K, et al. High glucose augments arginase activity and nitric oxide production in the renal cortex[J]. Metabolism, 2004, 53(7):868-874.

[3] Zharikov S, Krotova K, Hu H, et al. Uric acid decreases NO production and increases arginase activity in cultured pulmonary artery endothelial cells[J]. American Physiological Society, 2008(5).

Related Products:

- BC1500/BC1505 Plant Nitrate Nitrogen Assay Kit
- BC1520/BC1525 Plant Ammoniacal Nitrogen Assay Kit
- BC1530/BC1535 Urea Nitrogen (BUN) Assay Kit